

Mechanism for fractionation of Ta-Nb, Sn and W in granite-related metallogenic systems based on observations from the Karagwe-Ankole Belt of Central Africa

Niels HULSBOSCH¹, Stijn DEWAELE² & Philippe MUCHEZ¹

*1 Division of Geology, Department of Earth and Environmental Sciences, KU Leuven, Celestijnenlaan 200E, 3001 Leuven, Belgium. *niels.hulsbosch@kuleuven.be*

2 Mineralogy and Petrology, Department of Geology, Ghent University, Krijgslaan 281 S8, 9000 Ghent, Belgium.

Tantalum, niobium, tin and tungsten mineralization occurs worldwide in close proximity to fractionated peraluminous leucogranites. These granite-related ore deposits show a diversity of mineralization modes, ranging from plutonic-hosted breccia and hydrothermal vein-stockwork systems to (peri)batholithic greisens, skarns, pegmatites and hydrothermal veins. In the Mesoproterozoic Karagwe-Ankole orogenic belt (KAB) of Central-Africa numerous early-Neoproterozoic Nb-Ta-Sn-W rare-metal deposits formed as primary mineralization in hydrothermal quartz veins, magmatic lithium-cesium-tantalum family pegmatites and associated intra-pegmatitic greisens (Fig. 1). These deposits are on a belt-scale spatially and temporally related to ilmenite-series, peraluminous, S-type leucogranites, which are generally defined in the KAB as the Kibara Sn granite generation with an intrusion age of $\sim 986 \pm 10$ Ma (SHRIMP U-Pb on zircon; Tack et al., 2010).

This study reviews the petro- and metallogenesis of Nb-Ta-Sn-W granite-related ore deposits, as they occur in the Rwandese part of the KAB, and provides a general overview of the geochemical mechanisms behind the formation of pegmatite- and quartz vein-hosted deposits and the distribution and enrichment of the ore elements Nb, Ta, Sn and W in the different metallogenic subsystems (i.e. granites, pegmatites and veins). Moreover, the observed close spatiotemporally association between leucogranites, pegmatites and quartz veins is assessed in terms of a direct genetic link between the mineralization and the felsic magmatism. Based on this review and element distribution calculations, an integrated orthomagmatic metallogenic model for Nb-Ta-Sn-W mineralization in the KAB is demonstrated. Rayleigh-type fractional crystallization acted as the main mechanism by which pegmatitic magmas differentiate from a parental leucogranitic melt and by which incompatible elements Nb, Ta, Sn and W initially are enriched. However, early aqueous fluid immiscibility has been identified to occur during differentiation of this B-rich, F-poor melt system, which greatly effects W depletion in the melt by preferential partitioning of W into the mobile fluid phase. Sn retains dominantly in the melt phase and becomes, by high degrees of fractional crystallization, enriched in late-stage magmatic fluids. Element-specific melt-fluid-crystal fractionation together with element-specific and lithological-controlled precipitation conditions are the key enrichment processes and are all responsible for the decoupling of Nb-Ta, Sn and W and their subsequent precipitation in pegmatite, hydrothermal quartz vein and greisen deposits. This formation model for the KAB system can form a major tool in the exploration for granite-related ore deposits in general.




Figure 1. Conceptual outline of the geographical and geological setting of the major rare-metal deposits of the Karagwe-Ankole (KAB) and neighboring Kibara (KIB) belts in Central African Great Lakes region. B: Bujumbura, BB: Bengo-Biri, Bi: Bisesero, Bk: Bukavu, Bu: Busanga, Bug: Bugarama, GG: Gatumba-Gitarama pegmatite field, Gi: Gifurwe K: Kigali, Ka: Kampala, Kaa: Kabaya, Kab: Kabunga, Kai: Kailo, Kal: Kalanda, Kas: Kasese, Kl: Kalima, Ki: Kivuvu, Kib: Kibungo, KaM: Kamituga-Mobale, KM: Kirundo-Muyinga, KY: Kabambare – Yungwe, L: Lowa, Kis: Kiseke, Ma: Matala, Mi: Mitwaba, MK: Manono-Kitotolo, MN: Musha-Ntunga, Mu: Mulehe, Mua: Muana, Mui: Muika, Mw: Mwenga, N: Nyakabingo, Ny: Nyangulube P: Punia, R: Rutongo, Ru: Ruhembe, Rw: Rwinkwavu, S: Sasa. After Hulsbosch (in press).

References

- Hulsbosch, N, In press. Nb-Ta-Sn-W distribution in granite-related ore systems: fractionation mechanisms and examples from the Karagwe-Ankole Belt of Central Africa. In Decrée, S. & Robb, L. (eds), *Ore Deposits Origin Exploration and Exploitation*. Wiley. In press.
- Tack, L., Wingate, M.T.D., De Waele, B., Meert, J., Belousova, E., Griffin, B., Tahon, A. & Fernandez-Alonso, M., 2010. The 1375 Ma “Kibaran event” in Central Africa: Prominent emplacement of bimodal magmatism under extensional regime. *Precambrian Research*, 180, 63-84.